

This listing of claims will replace all prior versions and listings of claims in this application:

Listing of Claims

1. – 16. (Cancelled)

17. (Currently Amended) A method of making a bipathic medication, comprising the steps of:

making an active medicinal substance in a therapeutic dose from an initial substance;

using a homeopathic potentiation technology to make a potentiated medicinal preparation from the same initial substance; and

producing the bipathic medication by combining the active medicinal substance in the therapeutic dose and the potentiated medicinal preparation produced by the homeopathic potentiation technology, wherein the active medicinal substance and the potentiated medicinal preparation differ in their ~~being chemically homogeneous, but differing in a mechanism of~~ action on an organism and wherein the bipathic medication produced by combining the active medicinal substance and the potentiated preparation increases therapeutic efficiency of the active medicinal substance as compared to that of the active medicinal substance alone.

18. (Cancelled)

19. (Previously presented) The method as defined in claim 17, wherein said combining includes impregnating the active medicinal substance in the therapeutic dose with the potentiated medicinal preparation so that the active medicinal substance in the therapeutic dose is a carrier for the potentiated medicinal preparation.

20. (Previously presented) The method as defined in claim 17, wherein said combining includes dissolving the potentiated medicinal preparation in the active medicinal substance in the therapeutic dose so that the active medicinal substance in the therapeutic dose is a carrier for the potentiated medicinal preparation.

21. (Previously presented) The method as defined in claim 17, wherein said combining includes introducing the potentiated medicinal preparation into the active medicinal substance in the therapeutic dose formed as a paste so that the active medicinal substance in the therapeutic dose is a carrier for the potentiated medicinal preparation.

22. (Previously presented) The method as defined in claim 17, wherein said combining includes introducing the potentiated medicinal preparation method and the active medicinal substance in the therapeutic dose simultaneously into ~~an~~ the organism so that the active medicinal substance in the therapeutic dose and the potentiated medicinal preparation make said medication in the organism.

23. (Currently Amended) A bipathic medication, comprising
a therapeutic dose of an active medicinal substance; made from an initial substance
a potentiated medicinal preparation produced by a homeopathic potentiation method from the same initial substance, the potentiated medicinal preparation and being chemically homogeneous with the active medicinal substance differing in their action of an organism; and
a carrier having the same a chemical formula the same as the initial substance, the carrier comprising the active medicinal substance combined with the potentiated medicinal preparation;
wherein the bipathic medication produced by combining the active medicinal substance and the potentiated preparation increases therapeutic efficiency of the active medicinal substance as compared to that of the active medicinal substance alone.

24. (Cancelled)

25. (Previously presented) The medication as defined in claim 23, wherein said active medicinal substance in the therapeutic dose is impregnated with the potentiated medicinal preparation produced by the homeopathic potentiation method so that the

active medicinal substance in the therapeutic dose is a the carrier for the potentiated medicinal preparation.

26. (Previously presented) The medication as defined in claim 23, wherein said potentiated medicinal preparation produced by the homeopathic potentiation method is dissolved in the active medicinal substance in the therapeutic dose so that the active medicinal substance in the therapeutic dose is a the carrier for the potentiated preparation.

27. (Previously presented) The medication as defined in claim 23, wherein said potentiated medicinal preparation produced by the homeopathic potentiation method is introduced into the active medicinal substance in the therapeutic dose formed as a paste so that the active medicinal substance in the therapeutic dose is a the carrier for the potentiated medicinal preparation.

28. (Previously presented) The medication as defined in claim 23, wherein said potentiated medicinal preparation produced by the homeopathic potentiation method is introduced simultaneously into an organism with said active medicinal substance in the therapeutic dose, together making said medication in the organism.

29. (Previously presented) The medication as defined in claim 23, wherein said active medicinal substance in the therapeutic dose is atropine sulfate produced from atropine while said potentiated medicinal preparation produced by the homeopathic potentiation method is a C30 potency atropine sulfate dilution produced in accordance with the homeopathic potentiation method.

30. (Previously presented) The medication as defined in claim 23, wherein said active medicinal substance in the therapeutic dose is produced from acetylsalicylic acid, the potentiated medicinal preparation has a C30 potency and is produced by the homeopathic potentiation method from the acetylsalicylic acid, the medication being

obtained by impregnating milk sugar with the potentiated medicinal preparation having a C30 potency.

31. (Previously presented) The medication as defined in claim 23, wherein said active medicinal substance in the therapeutic dose includes 1.00 ml of prednizolon in the carrier impregnated with the potentiated medicinal preparation produced by the homeopathic potentiation method, comprising ~~is~~ 0.005 mg of potentiated prednizolon having a C12 potency.

32. (Previously presented) The medication as defined in claim 23, wherein said active medicinal substance in a therapeutic dose is insulin, and wherein said potentiated medicinal preparation produced by the homeopathic potentiation method is insulin having a C30 potency.

33. (Previously presented) The medication as defined in claim 23, wherein said active medicinal substance in the therapeutic dose is a paste from zinc, and wherein said potentiated medicinal preparation is produced by the homeopathic potentiation method from zinc.

34. (Previously presented) The medication as defined in claim 23, wherein said active medicinal substance in the therapeutic dose is sarcolysin dissolved in a potassium chloride, and wherein said potentiated medicinal preparation having a C200 potency is produced from the sarcolysin by the homeopathic potentiation method.

35. (Currently Amended) A method of ~~treatment~~ treating an organism with a medication, comprising the steps of:
making the medication comprising a carrier with therapeutic dose of an active medicinal substance made of an initial substance, the carrier having a chemical formula the same as the initial substance, and a potentiated medicinal preparation produced by a

homeopathic potentiation method from the same initial substance, wherein the active medicinal substance and the potentiated medicinal preparation differ in their action on the organism ~~and being chemically homogeneous with the active medicinal substance~~; and introducing the medication into ~~an~~ the organism;
wherein the bipathic medication produced by combining the active medicinal substance and the potentiated preparation increases therapeutic efficiency of the active medicinal substance as compared to that of the active medicinal substance alone.

36. (Currently Amended) The method of ~~treatment~~ treating as defined in claim 35, wherein making the medication comprises combining said active medicinal substance in the therapeutic dose and said potentiated medicinal preparation produced by the homeopathic potentiation method with one another to form the medication before introducing the medication into the organism.

36. (Currently Amended) The method of ~~treatment~~ treating as defined in claim 35, wherein the active medicinal substance in the therapeutic dose and the potentiated medicinal preparation produced by the homeopathic potentiation method are introduced into the organism separately; to be combined with one another in the organism.